

## Antibiotic Durations Quick Reference Table

Diagnosis	Recommended Duration	Source
<b>Acute Bacterial Rhinosinusitis (ABRS)</b>	5 days	<a href="#">IDSA Guidelines</a> (2012)
<b>Acute Otitis Media (AOM)</b>	<u>Age &lt; 2 years</u> : 10 days <u>Age ≥ 2 years</u> : 5 days	<a href="#">Cochrane Library</a> (2010) <a href="#">AAP Clinical Practice Guideline</a> (2013) <a href="#">CHCO Clinical Pathway</a> (2023)
<b>Appendicitis</b>	<u>Pre-Op</u> : One dose CTX and MTZ <u>Uncomplicated</u> : No additional abx <u>Complicated</u> : IV abx until clinically well or POD7. Obtain CBC on POD7 <ul style="list-style-type: none"> <li>• WBC ≤ 10: NO abx</li> <li>• WBC ≥ 10: 4 more days</li> </ul> <u>Abscess or phlegmon</u> : 8 days total from source control	<a href="#">CHCO Clinical Pathway</a> (2022)
<b><i>C. difficile</i> infection (CDI)</b>	10 days	<a href="#">CHCO Clinical Pathway</a> (2019)
<b>Group A Streptococcal (GAS) Pharyngitis</b>	10 days Note: consider shorter duration (1-5 days). See CHCO Pathway for more information.	<a href="#">IDSA Guidelines</a> (2012) <a href="#">CHCO Clinical Pathway</a> (2023)
<b>Uncomplicated Cystitis</b>	3 days	<a href="#">CHCO Clinical Pathway</a> (2021) <a href="#">IDSA Guidelines</a> (2010)
<b>Pyelonephritis</b>	7 days	<a href="#">CHCO Clinical Pathway</a> (2021) <a href="#">AAP Clinical Practice Guideline</a> (2011) <a href="#">IDSA Guidelines</a> (2010)
<b>Skin and Soft Tissue Infections (SSTI)</b>	5 days	<a href="#">IDSA Guidelines</a> (2014)
<b>Uncomplicated Community Acquired Pneumonia</b>	5 days	<a href="#">CHCO Clinical Pathway</a> (2022)
<b>Tracheitis associated with ET or tracheostomy tube</b>	5 days	<a href="#">Johns Hopkins Guidelines</a> (2011)
<b>Hospital-acquired and Ventilator associated pneumonia</b>	7 days	<a href="#">IDSA Guidelines</a> (2016)