

VENOUS THROMBOEMBOLISM (VTE) PREVENTION

ALGORITHM. Venous Thromboembolism (VTE) Prevention Inpatient




TABLE OF CONTENTS

[Algorithm. Venous Thromboembolism \(VTE\) Prevention Inpatient](#)

[Target Population](#)

[Background | Definitions](#)

[Initial Evaluation](#)

[Clinical Management](#)

[Parent | Caregiver Education](#)

[Key Contacts](#)

[References](#)

[Clinical Improvement Team](#)

TARGET POPULATION

Inclusion Criteria

Intended for:

- All patients with anticipated hospitalization greater than 23 hours

Exclusion Criteria

Not intended for:

- Patients with current VTE. Please consider Hematology consultation and refer to the Anticoagulation Dosing and Monitoring Protocol
- Patients in Neonatal Intensive Care Unit (NICU)
- Patients in the Labor and Delivery Unit
- Patients, 18 years and older, admitted to medical floors will be screened for VTE risk and prophylaxis ordered per the [Best Practice Advisory](#).
- Patients with moderate/severe Traumatic Brain Injury

BACKGROUND | DEFINITIONS

Definitions

VTE: Venous Thromboembolism, including deep vein thrombosis (of the limbs, abdomen, neck/chest, cranial), intracardiac thrombosis, and pulmonary embolism.

INITIAL EVALUATION

All patients should be assessed for VTE risk within 24 hours of admission and on a daily basis. Patients are determined to be “at-risk” depending on the risk factors present. Refer to the [Inpatient VTE algorithm](#).

VTE Risk Factors include:

- Age, greater than or equal to 12 years of age or post-pubertal
- Altered mobility: relative to patient's baseline, affecting one or more extremities greater than 48 hours (including that resulting from sedation for mechanical ventilation, acute spinal injury, transverse myelitis, or Guillain-Barre syndrome, etc.), defined as a Braden Q mobility score of 1 (completely immobile) or 2 (very immobile).
- Burns: greater than 50% of body surface area.
- Cancer: not in remission
- Cardiovascular low flow compromise: including, but not limited to, ventricular dysfunction or structural cardiac defect with associated turbulent blood flow (single ventricle physiology, cyanotic congenital heart defects, cavopulmonary anastomoses)
- Central venous catheter (CVC): includes peripherally inserted catheter (PICC), tunneled CVCs, non-tunneled CVCs, implanted ports, hemodialysis catheters
- COVID-19 infection, symptomatic requiring inpatient admission: Observational studies in adults have suggested that SARS-CoV-2 infection causes a prothrombotic state. The incidence of VTE in pediatric acute COVID-19 is unknown. For more information, refer to the [Guidelines for the Prevention and Treatment of VTE in Hospitalized Patients with COVID-19](#).
- Cystic fibrosis (CF)
- Estrogen therapy: currently taking (consider all formulations, including oral, transdermal patches, injections, etc.)
- Severe infection, confirmed (culture positive) or presumed requiring initiation of antimicrobials. Patients with osteomyelitis, particularly due to MRSA, are at high risk for developing VTE.
- Inflammatory disease flare, non-infectious: including, but not limited to, arthritis, inflammatory bowel disease (IBD), systemic lupus erythematosus (SLE), graft versus host disease (GVHD)
- Obesity: body mass index (BMI) greater than or equal to 95%-ile for gender/age (or greater than or equal to 30)
- Nephrotic syndrome, active
- Personal or 1st-degree family history of VTE or severe thrombophilia (acquired or inherited; however, new inpatient testing for purpose of risk categorization not recommended). For 1st-degree family history, only consider family members less than 50 years of age. For patients, consider whether they have been previously informed that they require blood thinners during hospitalization. Severe thrombophilia traits include:
 - Severe Antithrombin deficiency
 - Severe Protein S deficiency
 - Severe Protein C deficiency
 - Homozygous factor V Leiden mutation
 - Homozygous prothrombin mutation 20210
- Pregnancy: currently pregnant or within 6 weeks post-partum
 - Note: Low molecular weight heparin (LMWH) preferred over heparin for pregnant women due to FDA pregnancy category (does not cross placenta)
- Severe dehydration: including ongoing hyperosmolar acidotic state like diabetic ketoacidosis
- Surgery or procedure lasting longer than 90 minutes within the last 14 days
- Trauma: requiring an ICU admission. Excludes patients with acute traumatic brain injury. See the TBI: Moderate/Severe Clinical Pathway for more guidance

CLINICAL MANAGEMENT

Interventions

Appropriateness of therapeutic intervention is determined by each patient's level of risk balanced against any contraindications. In some circumstances, the patient may have multiple risk factors that are offset by clinical judgment, patient's activity level, family refusal of prophylaxis, etc. If a question remains about whether to initiate prophylaxis, consult hematology. A hematology consult is strongly recommended prior to starting pharmacologic prophylaxis outside of the ICU/CPCU.

General Measures

- Both dehydration and altered mobility are independent risk factors for thrombosis. Therefore, adequate hydration should be maintained and patients should be mobilized as early as possible.
- The primary care team should discuss the indications for all CVCs daily and should promptly remove any CVC that is no longer indicated.

Mechanical Prophylaxis

- Mechanical prophylaxis refers to either sequential compression devices (SCDs) or graduated compression stockings and is indicated for all patients at moderate-to-high risk for VTE, unless contraindicated.
- Mechanical prophylaxis can reduce clot formation through both physical and biochemical mechanisms. Compression of a blood vessel results in forward movement of blood and the shear and strain forces will trigger endothelium to release nitric oxide, prostacyclin, and tissue plasminogen activator. In addition, there is a reduction in plasminogen activator inhibitor.
- Contraindications to mechanical prophylaxis include:
 - Acute fracture of extremity (mechanical prophylaxis is to be applied to unaffected extremity)
 - Allergy to garment fabric
 - Osteogenesis imperfecta
 - Peripheral intravenous access in extremity (mechanical prophylaxis is to be applied to be applied to unaffected extremity)
 - Skin conditions affecting extremity mechanical prophylaxis is to be applied to (e.g., dermatitis, burns, recent skin grafts, leg wounds)
 - Suspected or existing deep vein thrombosis in extremity in question (can use graduated compression stockings in this circumstance)
 - Unable to achieve correct fit due to patient size
- The effectiveness of mechanical prophylaxis is dependent on the correct fit of the stocking or device. Patients should be measured to ensure proper fit.
- For patients that are unable to move on their own, passive range of motion (PROM) exercises should be completed with repositioning.
- Sequential Compression Devices (SCDs):
 - SCDs are available in adult and pediatric sizes for maximum calf sizes between 25 cm and 76 cm in circumference (no minimum circumference listed). Moleskin or other dermatologic protection may be necessary to prevent skin breakdown.

- Graduated compression stockings (if SCDs contraindicated, unavailable, or refused, or if patient has an existing DVT in that limb):
 - Graduated compression stockings reduce the overall cross-sectional area of the limb, increase linear velocity of venous flow, reduce venous wall distension, and improve valvular function.
 - Knee length stockings have been shown to be as effective as thigh length stockings and are associated with greater patient compliance and less skin breakdown.
 - Graduated compression stockings are manufactured for adults in small, medium, large, and extra-large sizes; regular and long lengths; light to firm support. Fit is determined according to measurements taken from thigh, calf, ankle, and foot. It is unlikely that graduated compression stockings will fit very young or small pediatric patients.

Pharmacologic prophylaxis

Note: Hematology consult is strongly recommended prior to starting pharmacologic prophylaxis outside of the ICU/CPCU, since there is lack of evidence demonstrating the safety or efficacy of pharmacologic prophylaxis of VTE in children.

Patients under the care of a surgeon must have approval from the attending surgeon prior to initiation of pharmacologic prophylaxis.

Pharmacologic prophylaxis, most commonly with low-molecular weight heparin (enoxaparin) or unfractionated heparin, should be considered for high-risk patients without contraindications. Risks commonly associated with any pharmacological prophylaxis include bleeding and bruising at injection site but could also include severe hemorrhagic complications (although this is unlikely at prophylactic dosing).

Clinical and possible laboratory monitoring of patients receiving pharmacologic prophylaxis:

- Patients should be observed for signs/symptoms of bleeding and therapy should be discontinued if bleeding is clinically significant.
- Low-molecular weight heparin may be held twelve (12) hours prior to any invasive procedure.
- In general, laboratory monitoring of anti-Xa levels (heparin assay) is not warranted. There are, however, certain circumstances in which a patient with expected alterations in pharmacokinetics (e.g., obesity, renal insufficiency, acute thermal burns and states of high inflammation) should have a heparin assay obtained to validate appropriateness of prophylactic dosing.
- If checking levels is warranted, typical VTE prophylaxis target for enoxaparin is between 0.2-0.4 units/mL (collected 4-hrs post-dose, using the heparin, low-molecular weight assay). Typical VTE prophylaxis target for unfractionated heparin is between 0.1-0.3 units/mL (collected 4-hrs after every dose change, using the heparin, unfractionated assay).

Contraindications to pharmacologic prophylaxis include:

- Absolute:
 - Evidence of active hemorrhage
 - Epidural catheter or lumbar puncture performed within last hour^[OBJ]
 - Intravascular thrombolytic therapy within the last 24 hours
 - Known arterio-venous malformation, aneurysm, or moyamoya
 - Acute period following neurosurgery or severe traumatic brain injury (see appendix B in the [TBI Moderate/Severe Clinical Pathway](#) for more guidance)

- Acute large-territory arterial ischemic stroke
- Platelet count unable to be sustained above 30,000/mm³ (platelet transfusions for the sole purpose of elevating the platelet count to an acceptable level for pharmacologic prophylaxis are not recommended)
- Plasma fibrinogen concentration unable to be maintained above 100 g/dL
- Prior history of unexplained spontaneous hemorrhage
- For Low Molecular Weight Heparin (LMWH): known allergy/anaphylaxis to LMWH⁽⁰⁹⁾ or pork products
- Relative:
 - Known bleeding disorder or hemophilia
 - Recent hemorrhage
 - Uncontrolled hypertension
 - Unexplained coagulopathy
 - Liver failure-associated coagulopathy
 - For Low Molecular Weight Heparin (LMWH): moderate or severe renal insufficiency, with creatinine clearance less than 30 mL/min

Dosing:

- See CHCO Formulary for dosing:
- [Low molecular weight heparin \(LMWH\)/enoxaparin](#)
- [Unfractionated heparin \(uFH\)](#)

PARENT | CAREGIVER EDUCATION

Provide parents and caregivers with information regarding VTE prophylaxis, the need for it, and any risks associated with the indicated prophylaxis.

KEY CONTACTS

Consult with Anticoagulation Clinical Pharmacist for specific questions around pharmacologic prophylaxis.

REFERENCES

1. Sandoval JA, Sheehan MP, Stonerock CE, Shafique S, Rescorla FJ, Dalsing MC. Incidence, risk factors, and treatment patterns for deep venous thrombosis in hospitalized children: an increasing population at risk. *J Vasc Surg* 2008;47:837-43.
2. Hanson SJ, Punzalan RC, Arca MJ, et al. Effectiveness of clinical guidelines for deep vein thrombosis prophylaxis in reducing the incidence of venous thromboembolism in critically ill children after trauma. *J Trauma Acute Care Surg* 2012;72:1292-7.
3. Vu LT, Nobuhara KK, Lee H, Farmer DL. Determination of risk factors for deep venous thrombosis in hospitalized children. *J Pediatr Surg* 2008;43:1095-9.
4. Branchford BR, Mourani P, Bajaj L, Manco-Johnson M, Wang M, Goldenberg NA. Risk factors for in-hospital venous thromboembolism in children: a case-control study employing diagnostic validation. *Haematologica* 2012;97:509-15.
5. Caruso V, Iacoviello L, Di Castelnuovo A, et al. Thrombotic complications in childhood acute lymphoblastic leukemia: a meta-analysis of 17 prospective studies comprising 1752 pediatric patients. *Blood* 2006;108:2216-22.
6. Walker AJ, Grainge MJ, Card TR, West J, Ranta S, Ludvigsson JF. Venous thromboembolism in children with cancer - A population-based cohort study. *Thromb Res* 2013.
7. Sharathkumar AA, Mahajerin A, Heidt L, et al. Risk-prediction tool for identifying hospitalized children with a predisposition for development of venous thromboembolism: Peds-Clot clinical Decision Rule. *J Thromb Haemost* 2012;10:1326-34.
8. Higgerson RA, Lawson KA, Christie LM, et al. Incidence and risk factors associated with venous thrombotic events in pediatric intensive care unit patients. *Pediatr Crit Care Med* 2011;12:628-34.
9. Hanson SJ, Punzalan RC, Greenup RA, Liu H, Sato TT, Havens PL. Incidence and risk factors for venous thromboembolism in critically ill children after trauma. *J Trauma* 2010;68:52-6.
10. Chalmers EA. Epidemiology of venous thromboembolism in neonates and children. *Thromb Res* 2006;118:3-12.
11. Pomp ER, Lenselink AM, Rosendaal FR, Doggen CJ. Pregnancy, the postpartum period and prothrombotic defects: risk of venous thrombosis in the MEGA study. *J Thromb Haemost* 2008;6:632-7.
12. Goldman RD, Friedman JN, Parkin PC. Validation of the clinical dehydration scale for children with acute gastroenteritis. *Pediatrics* 2008;122:545-9.
13. Covidien. T.E.D. Anti-Embolism Stockings. In. Mansfield; 2010.
14. Sajid MS, Desai M, Morris RW, Hamilton G. Knee length versus thigh length graduated compression stockings for prevention of deep vein thrombosis in postoperative surgical patients. *Cochrane Database Syst Rev* 2012;5:CD007162.
15. Horlocker TT, Wedel DJ, Rowlingson JC, et al. Regional anesthesia in the patient receiving antithrombotic or thrombolytic therapy: American Society of Regional Anesthesia and Pain Medicine Evidence-Based Guidelines (Third Edition). *Reg Anesth Pain Med* 2010;35:64-101.
16. Lewis TV, Johnson PN, Nebbia AM, Dunlap M. Increased enoxaparin dosing is required for obese children. *Pediatrics* 2011;127:e787-90.
17. Urbankova et al. Intermittent pneumatic compression and deep vein thrombosis prevention: a meta-analysis in postoperative patients. *Thromb Haemost* 2005;94:1191-5.
18. Massicotte et al. PROTEKT Study Group. An open label randomized controlled trial of low molecular weight heparin for the prevention of central venous line-related thrombotic complications in children; the PROTEKT trial. *Thromb Res* 2003;109(2-3):101-108.
19. Reiter PD, Wathen B, Valuck RJ, Dobyens EL. Thrombosis risk factor assessment and implication for prevention in critically ill children. *Pediatr Crit Care Med* 2012 Jul; 13(4):381-6.
20. Monagle P et al. Antithrombotic therapy in neonates and children: antithrombotic therapy and prevention of thrombosis, 9 ed: American College of Chest Physicians evidence-based clinical practice guidelines. *Chest* 2012;141:e737S-e801S.
21. Stem J et al. Safety of prophylactic anticoagulation at a pediatric hospital. *J Pediatr Hematol Oncol*. 2013; 35(7):e287-291.
22. Hofman S et al. Clinical experience with low-molecular weight heparins in pediatric patients. *Thromb Res* 2001;103(5):345-53.

23. Dix D et al. The use of low molecular weight heparin in pediatric patients: a prospective cohort study. *J Pediatr* 2000;136:439-45
24. Bidlingmair C et al. Safety and efficacy of low molecular weight heparins in children: a systematic review of the literature and meta-analysis of single-arm studies. *Sem Thromb Hemos* 2011;37(7):814-25.
25. Amankwah, EK et al. Risk factors for hospital-associated venous thromboembolism in the neonatal care unit. *Thromb Res* 2014;134(2):305-9.
26. Atchison CM et al. Development of a new risk score for hospital-acquired venous thromboembolism in noncritically ill children: findings from a large single-institutional case control study. *J Pediatr* 2014;165(4):793-8.
27. Arlikar SJ et al. Development of a new risk score for hospital-acquired venous thromboembolism in critically-ill children not undergoing cardiothoracic surgery. *Thromb Res* 2015;136(4):717-22.
28. Mahajerin A et al. Hospital-acquired venous thromboembolism in pediatrics: a systematic review and metaanalysis of risk factors and risk-associated models. *Haematologica* 2015;100(8):1045-50.
29. Hanson SJ et al. Recommendations for venous thromboembolism prophylaxis in pediatric trauma patients: a national, multidisciplinary consensus study. *J Trauma Acute Care Surg* 2016.
30. Al-Samkari H, Karp Leaf RS, Dzik WH, Carlson JCT, Fogerty AE, Waheed A, Goodarzi K, Bendapudi PK, Bornikova L, Gupta S, Leaf DE, Kuter DJ, Rosovsky RP. COVID-19 and coagulation: bleeding and thrombotic manifestations of SARS-CoV-2 infection. *Blood*, 2020 136: 489–500



CLINICAL IMPROVEMENT TEAM MEMBERS

Michele Loi, MD | Critical Care and CCBD
John S. Kim, MD, MS | Cardiology and Cardiac ICU
Susan Kuldaneck, MD | Hematology
Kelly Reichert, MS, FNP-C, CCRN | Clinical Practice Specialist
Michael Wang, MD | Hematology

APPROVED BY

Todd Carpenter, MD | Critical Care
Pamela D Reiter, PharmD | Clinical Pharmacy Specialist, Pediatric ICU
Timothy Schardt, PharmD, BCPS | Clinical Pharmacist for Investigational Drug Services and Anticoagulation

Clinical Pathways and Measures Committee – April 26, 2021
 Venous Thromboembolism Steering Committee – January 26, 2021
 Pharmacy & Therapeutics – May 6, 2021

MANUAL/DEPARTMENT	Clinical Care Guidelines/Quality
ORIGINATION DATE	October 4, 2011
LAST DATE OF REVIEW OR REVISION	May 21, 2021
COLORADO SPRINGS REVIEWED BY	 Michael DiStefano, MD Chief Medical Office, Colorado Springs
APPROVED BY	 Lalit Bajaj, MD Chief Quality & Outcomes Officer, Clinical Effectiveness

REVIEW | REVISION SCHEDULE

Scheduled for full review on May 6, 2025

Clinical pathways are intended for informational purposes only. They are current at the date of publication and are reviewed on a regular basis to align with the best available evidence. Some information and links may not be available to external viewers. External viewers are encouraged to consult other available sources if needed to confirm and supplement the content presented in the clinical pathways. Clinical pathways are not intended to take the place of a physician's or other health care provider's advice, and is not intended to diagnose, treat, cure or prevent any disease or other medical condition. The information should not be used in place of a visit, call, consultation or advice of a physician or other health care provider. Furthermore, the information is provided for use solely at your own risk. CHCO accepts no liability for the content, or for the consequences of any actions taken on the basis of the information provided. The information provided to you and the actions taken thereof are provided on an "as is" basis without any warranty of any kind, express or implied, from CHCO. CHCO declares no affiliation, sponsorship, nor any partnerships with any listed organization, or its respective directors, officers, employees, agents, contractors, affiliates, and representatives.

Discrimination is Against the Law. Children's Hospital Colorado complies with applicable Federal civil rights laws and does not discriminate on the basis of race, color, national origin, age, disability, or sex. Children's Hospital Colorado does not exclude people or treat them differently because of race, color, national origin, age, disability, or sex.

Children's Hospital Colorado provides free aids and services to people with disabilities to communicate effectively with us, such as: Qualified sign language interpreters, written information in other formats (large print, audio, accessible electronic formats, other formats). Children's Hospital Colorado provides free language services to people whose primary language is not English, such as: Qualified interpreters, information written in other languages.

If you need these services, contact the Medical Interpreters Department at 720.777.9800.

If you believe that Children's Hospital Colorado has failed to provide these services or discriminated in another way on the basis of race, color, national origin, age, disability, or sex, you can file a grievance with: Corporate Compliance Officer, 13123 E 16th Avenue, B450, Aurora, Colorado 80045, Phone: 720.777.1234, Fax: 720.777.7257, corporate.compliance@childrenscolorado.org. You can file a grievance in person or by mail, fax, or email. If you need help filing a grievance, the Corporate Compliance Officer is available to help you.

You can also file a civil rights complaint with the U.S. Department of Health and Human Services, Office for Civil Rights, electronically through the Office for Civil Rights Complaint Portal, available at ocrportal.hhs.gov/ocr/portal/lobby.jsf, or by mail or phone at: U.S. Department of Health and Human Services 200 Independence Avenue, SW Room 509F, HHH Building Washington, D.C. 20201 1-800-368-1019, 800-537-7697 (TDD) Complaint forms are available at www.hhs.gov/ocr/office/file/index.html.

Children's Hospital Colorado complies with applicable Federal civil rights laws and does not discriminate on the basis of race, color, national origin, age, disability, or sex.

ATENCIÓN: si habla español, tiene a su disposición servicios gratuitos de asistencia lingüística. Llame al 1-720-777-9800.

CHÚ Ý: Nếu bạn nói Tiếng Việt, có các dịch vụ hỗ trợ ngôn ngữ miễn phí dành cho bạn. Gọi số 1-720-777-9800.

주의: 한국어를 사용하시는 경우, 언어 지원 서비스를 무료로 이용하실 수 있습니다. 1-720-777-9800 번으로 전화해 주십시오.

注意: 如果您使用繁體中文, 您可以免費獲得語言援助服務。請致電1-720-777-9800。

ВНИМАНИЕ: Если вы говорите на русском языке, то вам доступны бесплатные услуги перевода. Звоните 1-720-777-9800.

ማስታወሻ: የሚናገሩት ቋንቋ አማርኛ ከሆነ የትርጉም አገዳታ ኗርጁት: በነጻ ሊያገዝዎት ተዘጋጅተዋል: ወደ ሚከተለው ቁጥር ይደውሉ 1-720-777-9800 (መስማት ለተሳናቸው).

ملحوظة: إذا كنت تتحدث انكر اللغة، فإن خدمات المساعدة اللغوية تتوفر لك بالمجان. اتصل برقم 720-777-9800-1 (رقم)

ACHTUNG: Wenn Sie Deutsch sprechen, stehen Ihnen kostenlos sprachliche Hilfsdienstleistungen zur Verfügung. Rufnummer: 1-720-777-9800.

ATTENTION : Si vous parlez français, des services d'aide linguistique vous sont proposés gratuitement. Appelez le 1-720-777-9800.

ध्यान बनु होस्तपाइले नेपाल बोलनहनछ भन तपाइको निम्त भाषा सहायता सवाहूर नःशल्क रूपमा उपलब्ध छ । फोन गनु होसरू 1-720-777-9800 ।

PAUNAWA: Kung nagsasalita ka ng Tagalog, maaari kang gumamit ng mga serbisyo ng tulong sa wika nang walang bayad. Tumawag sa 1-720-777-9800.

注意事項: 日本語を話される場合、無料の言語支援をご利用いただけます。1-720-777-9800 まで、お電話にてご連絡ください。

Nti: Q buri na asu lbo, asusu aka qasu n'efu, defu, aka. Call 1-720-777-9800.